

Reshaping-induced spatiotemporal chaos in driven, damped sine-Gordon systems

R. Chacón

*Departamento de Electrónica e Ingeniería Electromecánica,
Escuela de Ingenierías Industriales, Universidad de Extremadura, E-06071*

Badajoz, Spain

(February 6, 2008)

Typeset using REVTeX

Abstract

Spatiotemporal chaos arising from the competition between sine-Gordon-breather and kink-antikink-pair solitons by reshaping an ac force is demonstrated. After introducing soliton collective coordinates, Melnikov's method is applied to the resulting effective equation of motion to estimate the parameter-space regions of the ac force where homoclinic bifurcations are induced. The analysis reveals that the chaos-order threshold exhibits sensitivity to small changes in the force shape. Computer simulations of the sine-Gordon system show good agreement with these theoretical predictions.

I. INTRODUCTION

Although nonlinearity is an ubiquitous and fascinating feature of real-world dynamical phenomena, some aspects of such a fundamental property remain poorly explored, partly because they are rarely incorporated into the mathematical models which aim to describe even the simplest of such phenomena. In the context of nonautonomous systems, one of these aspects is the nature of the periodic excitations which can act on a given nonlinear, dissipative or Hamiltonian, system. Since nonlinear systems represent the general case, it would be natural to model periodic excitations (external or parametric) by using periodic functions that are solutions of (physically relevant) nonlinear equations, instead of harmonic functions which are solutions of linear systems, even though the latter have been overwhelmingly used up to now. To consider nonlinear periodic excitations implies enlarging the amplitude-period parameter space to include the parameters, α_i , that control the excitation shape. Note that, in the general case, these parameters α_i are the (normalized) Fourier coefficients of the periodic function. In physical terms this means that, for a fixed amplitude and period, the parameters α_i are responsible for the temporal rate at which energy is transferred from the excitation mechanism to the system. In the context of systems described by ordinary differential equations, this idea has led to the demonstration of the existence of new generic

temporal chaos $\leftrightarrow$ order routes by changing only the shape of a nonlinear periodic excitation [1-3].

In this present work, the onset of *spatio-temporal* chaos associated with the competition between sine-Gordon-breathers and kink-antikink-pairs is investigated in a one-dimensional sine-Gordon (sG) system

$$\Phi_{tt} - \Phi_{xx} + \sin \Phi = -\delta \Phi_t + \gamma \operatorname{sn} [4K(m)t/T; m], \quad (1)$$

when *only* the excitation shape is varied. Here, $\operatorname{sn}(u; m)$ is the Jacobian elliptic function (JEF) of parameter m ($K(m)$ is the complete elliptic integral of the first kind) used *solely* as an example to illustrate the reshaping-induced order-chaos route. When $m = 0$, then $\operatorname{sn} [4K(m = 0)t/T; m = 0] = \sin(2\pi t/T)$, i.e., one recovers the previously studied case of harmonic excitation [4]. This is relevant in comparing the structural stability of the sG system when solely the excitation shape is varied. In the other limit, $m = 1$, one obtains the Fourier expansion of the square wave function: $\operatorname{sn} [4K(m = 1)t/T; m = 1] = (4/\pi) \sum_{n=0}^{\infty} (2n + 1)^{-1} \sin [(2n + 1) 2\pi t/T]$. From an experimental standpoint, the choice of the above JEF is motivated for the fact that only the excitation shape is varied by increasing m from 0 to 1, and there is thus a smooth transition from a sine function to a square wave. While these two limiting shapes are readily implemented in experiments, no theoretical analysis connecting them has been developed up to now in the context of sG systems.

The organization of the paper is as follows. In Section 2, a collective coordinate approach [4] is used to obtain an effective equation of motion for the collective variables. In Section 3, Melnikov's method is applied to the resulting effective equation of motion to estimate the parameter-space regions of the elliptic ac force where the onset of chaotic instabilities is expected. Numerical evidence supporting the theoretical predictions from previous sections is presented in Section 4. Finally, Section 5 includes a summary of the findings.

II. COLLECTIVE COORDINATES

In this present work, a single-breather excitation is assumed to be present in the absence of dissipation and forcing ($\delta = \gamma = 0$). It is well known that the simultaneous action of dissipative ($\delta > 0$) and spatially uniform harmonic forces ($\gamma \sin(2\pi t/T)$) can break the breather into a kink-antikink pair, which can then recombine into a breather soliton [4,5]. For fixed dissipation, this process may occur repeatedly yielding an intermittent chaotic sequence for certain values of amplitude and period. The above elliptic ac driving permits one to study such a competition between the breather and the kink-antikink pair for parameter values belonging to *larger* regions of the amplitude-period parameter space. To this end, one makes a severe mode truncation to the breather collective coordinates [6,7], which yields a set of ordinary differential equations capable of being studied with the aid of Melnikov's method [8,9]. Such an effective equation exhibits a separatrix dynamics which is equivalent to the separatrix dynamics of the full sine-Gordon equation. In particular, homoclinic bifurcations correspond to (chaotic) intermittent binding and unbinding of the kink-antikink pair. In order to determine the equations of motion for the breather collective modes, one first needs the breather and kink-antikink solutions to the unperturbed sine-Gordon equation ($\delta = \gamma = 0$ in Eq. (1)) [4].

Breather:

$$\begin{aligned} U_B(x, t) &\equiv 4 \tan^{-1} \left\{ \frac{\tan \theta \sin [v(t)]}{\cosh [2k(x - x_0)]} \right\}, \\ v(t) &\equiv v_0 + t \cos \theta, \\ k &\equiv \frac{1}{2} \sin \theta, \\ E &\equiv 16 \sin \theta. \end{aligned} \tag{2}$$

Kink-antikink pair ($K \bar{K}$):

$$\begin{aligned} U_{K\bar{K}}(x, t) &\equiv 4 \tan^{-1} \left\{ \frac{\sinh [u(t - t_0)/(1 - u^2)^{1/2}]}{u \cosh [(x - x_0)/(1 - u^2)^{1/2}]} \right\}, \\ E_{K\bar{K}} &\equiv \frac{16}{(1 - u^2)^{1/2}} > 16, u \geq 0. \end{aligned} \tag{3}$$

Here, u is the $t \rightarrow \infty$ velocity of the kink while $-u$ is the $t \rightarrow \infty$ velocity of the antikink, and v_0, x_0, θ ($0 \leq \theta \leq 2\pi$) are constants. Also, $\omega_B \equiv \cos \theta$ is the breather's internal frequency and $E, E_{K\bar{K}}$ are the breather and kink-antikink-pair energies, respectively. Following Ref. [4], one assumes the *ansatz* that the solution of the perturbed sine-Gordon equation has the same form as the unperturbed breather (2), but now allowing θ and v_0 to be temporal functions:

$$v(t) = v_0(t) + \int^t \cos \theta(t') dt'. \quad (4)$$

This approximation means that one is assuming a perturbation sufficiently weak so that its principal effect on the system is to continuously alter the frequency and the phase of the breather. Therefore, such an approximation is valid when $\omega_B \ll 1$ since the breather energy is $E_B \equiv 16(1 - \omega_B^2)^{1/2}$. Next, one can deduce the equations of motion for the breather collective modes (u, v) , where

$$\begin{aligned} u(t) &\equiv U_B(x = x_0, t) = 4 \tan^{-1} A(t), \\ v(t) &\equiv \left[\frac{\partial U_B}{\partial t} \right]_{x=x_0} = \frac{4dA(t)/dt}{1 + A(t)^2}, \\ A(t) &\equiv \tan \theta \sin v(t). \end{aligned} \quad (5)$$

After the coordinate transformation $z \equiv \tan(u/4)$, $\dot{z} \equiv w$, such equations of motion can be shown to be

$$\begin{aligned} \dot{\chi} &= \mathbf{f}(\chi) + \epsilon \mathbf{g}(\chi, \psi), \\ \dot{\psi} &= 4K(m)/T, \end{aligned} \quad (6)$$

with

$$\begin{aligned} \chi &= \begin{pmatrix} z \\ w \end{pmatrix} \\ \mathbf{f}(\chi) &= \begin{pmatrix} w \\ -z(1 - w^2)(1 + z^2)^{-1} \end{pmatrix}, \end{aligned} \quad (7)$$

and the perturbation

$$\epsilon \mathbf{g}(\chi; t) = \begin{pmatrix} 0 \\ h(t)g_p(z, w) - \delta g_\delta(z, w) \end{pmatrix}, \quad (8)$$

where

$$\begin{aligned} h(t) &= \frac{\pi}{4} \gamma \operatorname{sn} [4K(m)t/T; m], \\ g_p(z, w) &= \frac{[(w^2 + z^2)z^2 - s\{z^4 + w^2[z^2 + (1 - w^2)(1 + z^2)]]}{(1 + sw^2)(w^2 + z^2)(1 + s)(1 + z^2)^{1/2}} \\ &\quad + \frac{2w^2z^2s^2(1 + z^2) + w^2s^3(1 + 2z^2)(1 + z^2)^2}{(1 + sw^2)(w^2 + z^2)(1 + s)(1 + z^2)^{1/2}}, \\ g_\delta(z, w) &= \frac{w[(2 + z^2)(w^2 + z^2)z^2 - s\{z^4 - w^2[(2 + z^2)z^4 - (1 - w^2)(1 + z^2)]]}{(1 + sw^2)(w^2 + z^2)(1 + z^2)} \\ &\quad + \frac{w[w^2s^2(1 + 2z^2)(1 + z^2)]}{(1 + sw^2)(w^2 + z^2)(1 + z^2)}, \\ s &\equiv \frac{\sinh^{-1}(z)}{z(1 + z^2)^{1/2}}. \end{aligned} \quad (9)$$

Noting that the separatrix in the phase-plane $u - v$ is given by

$$\begin{aligned} u_{0,\pm}(t) &= \pm 4 \tan^{-1}(t), \\ v_{0,\pm}(t) &= \pm 4/(1 + t^2), \end{aligned} \quad (10)$$

one finds that on the separatrix in the phase-plane $z - w$

$$\begin{aligned} z_{0,\pm}(t) &= \pm t, \\ w_{0,\pm}(t) &= \pm 1, \end{aligned} \quad (11)$$

where the positive (negative) sign refers to the upper (lower) homoclinic orbit in the phase-plane $u - v$, which is obtained for the energy $E = 16$ of the unperturbed energy functional $E = 2 \tan^2(u/4) + (u/4)^2 \sec^2(u/4)$.

III. CHAOTIC THRESHOLD FUNCTION

Now one can estimate the threshold of chaotic instabilities for the effective system (6) using Melnikov's method. Thus, the corresponding Melnikov function is

$$M^\pm(t_0) = \mp D + C \sum_{n=0}^{\infty} a_n(m) p_n(T) \sin \left[\frac{(2n+1) 2\pi t_0}{T} \right], \quad (12)$$

where

$$\begin{aligned} D &\equiv \delta \int_{-\infty}^{\infty} \frac{(1 - \tanh^2 x) [(2 - \tanh^2 x) \tanh^2 x + r + r^2 (1 - \tanh^4 x)]}{1 + r} dx \simeq 2.03727\delta, \\ p(2\pi/T) &\equiv \int_{-\infty}^{\infty} \frac{\cos\left(\frac{2\pi}{T} \sinh x\right) [(1-r)(1 - \tanh^2 x)] \operatorname{sech} x + 2r^2 \tanh^2 x + r^3 (1 + \tanh^2 x)}{(1+r)^2} dx, \\ C &\equiv \pi^2 \gamma / (4\sqrt{m}K(m)), \\ a_n(m) &\equiv \operatorname{csch} [(2n+1) \pi K(1-m)/(2K(m))], \\ p_n(T) &\equiv p[(2n+1) 2\pi/T], \end{aligned} \quad (13)$$

and where $r \equiv 2x \operatorname{csch}(2x)$. From Eq. (12) one sees that a homoclinic bifurcation is guaranteed if

$$\frac{\delta}{\gamma} < U(m, T), \quad (14)$$

where the chaotic threshold function is

$$U(m, T) \equiv \frac{\pi^2}{4\alpha\sqrt{m}K(m)} \sum_{n=0}^{\infty} (-1)^n a_n(m) p_n(T). \quad (15)$$

The series in Eq. (15) satisfies the conditions of the Leibnitz theorem on alternate series, and then one can deduce a slightly more restrictive condition for the onset of the aforementioned spatio-temporal chaotic instabilities, in order to facilitate the comparison with the previously studied case [4] of a harmonic force ($m = 0$):

$$\frac{\delta}{\gamma} < U(m = 0, T) R(m), \quad (16)$$

where $U(m = 0, T) = \pi p(2\pi/T)/(4\alpha)$, $R(m) \equiv \pi (\sqrt{m}K(m))^{-1} \operatorname{csch} [\pi K(1-m)/(2K(m))]$.

It is worth noting that the condition (16) is a reliable approximation due to the coefficients $a_n(m) \simeq 0$, $\forall n \geq 1$ and $\forall m \in [0, m^*]$ with m^* being very close to 1. A plot of the shape function $R(m)$ is shown in Fig. 1, where one sees that $R(m = 0) = 1$, as expected. The

rapid growth of $R(m)$ as $m \rightarrow 1$ means that the chaotic threshold exhibits hypersensitivity to changes in the excitation shape as this approximates a square wave. This can be understood in terms of the increasing accumulation of *effective* harmonics in the Fourier series of the elliptic force as $m \rightarrow 1$. Note that this number of effective harmonics is not a linear function of m because of the dependence of the elliptic force on $K(m)$.

IV. NUMERICAL SIMULATIONS

For the purpose of the comparison with the above theoretical predictions, the parameters $\delta = 0.2$, $T = 2\pi/0.6$, and $L = 24$ were fixed, and an sG breather with frequency $\omega_B = \sqrt{2}/2$ and $x_0 = v_0 = 0$ (cf. Eq. (2)) was chosen as initial data. The period $T = 2\pi/0.6$ corresponds to a frequency $\omega = 0.6$ (for a sinusoidal force, $m = 0$) for which the chaotic attractors are dominated by breather to kink-antikink transitions [11,12]. Equation (1) was integrated under periodic boundary conditions $\Phi(x = -L/2, t) = \Phi(x = L/2, t)$ for all t . Figure 2 shows an illustrative example of the comparison between the theoretically obtained chaotic threshold amplitude γ_{th} (solid line, cf. Eq. (16)) and the corresponding numerically obtained amplitude (dots) as the shape parameter m varies. One sees that the amplitudes obtained numerically are slightly larger than the respective theoretically predicted ones, as expected, although an overall good agreement is found. Similar good agreement was found for other sets of parameters in the large-period regime ($T \gg 2\pi$), provided that the damping and driving force were sufficiently weak. An illustrative sequence of the order-chaos route is shown in Fig. 3 where the initial state ($m = 0$) occurs after the system has undergone a spatial period doubling (for $m = 0$ and $0.6 \lesssim \gamma \lesssim 0.9$, cf. Ref. [12]) but remains simply periodic in time. This two-breather state has thus a spatial period $\frac{1}{2}$ and a temporal period 1, and exhibits mirror symmetry with respect to the origin. The spatial symmetry is broken as the forcing shape deviates from sinusoidal. Figure 3 illustrates this phenomenon with the values $m = 0.8$ and $m = 0.82$. This spatial symmetry breaking of the two-breather state appears to be an unavoidable precursor of the onset of spatio-temporal chaos via the

kink-antikink collision mechanism, as can be appreciated in Fig. 3 for the shape parameter $m = 0.898 \simeq m_{th}$, which corresponds to the theoretically predicted chaotic threshold (cf. Eq. (16)). For higher values of the shape parameter, i.e., when the driving shape is ever closer to a square wave, one finds the persistence of the chaotic dynamics (cf. Figs. 3, $m = 0.99$) predicted by the Melnikov-method-based analysis.

V. CONCLUSIONS

In sum, it has been shown through the example of an elliptic ac driving that the competition between sine-Gordon-breather and kink-antikink-pair solitons strongly depends on the shape of the spatially uniform ac driving. In particular, it was theoretically predicted and numerically confirmed that a square wave is more effective than a sinusoidal function to yield spatio-temporal chaos associated with the competition between sine-Gordon-breathers and kink-antikink-pairs. Potential application of the present results to reshaping-induced control (suppression and enhancement) of chaotic dynamics [1,14,15] range from DNA molecules [16] to long Josephson junctions and quasi-one-dimensional ferromagnets [17]. An interesting problem under current investigation is whether the above predictions on the chaotic threshold remain reliable in the so-called (for $m = 0$) nonlinear (cubic) Schrödinger regime (i.e., $T \lesssim 2\pi$), where the chaotic dynamics is dominated by breather-breather transitions instead of the breather to kink-antikink transitions of the regime studied here ($T \gg 2\pi$), since both mechanisms involve homoclinic crossings [13].

Acknowledgement 1 *The author acknowledges financial support from the Spanish MCyT and the European Regional Development Fund (FEDER) program through project FIS2004-02475.*

REFERENCES

- [1] Chacón R, Díaz Bejarano J. Routes to suppressing chaos by weak periodic perturbations. *Phys Rev Lett* 1993; 71: 3103-3106.
- [2] Zeni AR, Gallas JAC. Lyapunov exponents for a Duffing oscillator. *Physica D* 1995; 89: 71-82.
- [3] Chacón R, Martínez García-Hoz A. Route to chaos via strange non-chaotic attractors by reshaping periodic excitations. *Europhys Lett* 2002; 57: 7-13.
- [4] Ariyasu JC, Bishop AR. Determination of the onset of spatiotemporal complexity induced by soliton competition in a perturbed sine-Gordon chain. *Phys Rev A* 1989; 39: 6409-6415.
- [5] Bishop AR, McLaughlin DW, Salerno M. Global coordinates for the breather-kink (antikink) sine-Gordon phase space: An explicit separatrix as a possible source of chaos. *Phys Rev A* 1989; 40: 6463-6469.
- [6] Ercolani NM, Forest MG, McLaughlin DW, Montgomery R. Hamiltonian structure of the modulation equations for sine-Gordon wavetrains. *Duke Math J* 1987; 55: 949-983.
- [7] Salerno M. Reduced sine-Gordon breather-(anti)kink dynamics and the double sine-Gordon system. *Phys Lett A* 1989; 134: 421-423.
- [8] Melnikov VK. On the stability of the center for time periodic perturbations. *Trans Moscow Math Soc* 1963; 12: 1-57.
- [9] Guckenheimer J, Holmes PJ. Nonlinear oscillations, dynamical systems, and bifurcations of vector fields. New York: Springer; 1983.
- [10] Bishop AR, Fesser K, Lomdahl PS, Kerr WC, Williams MB, Trullinger SE. *Phys Rev Lett* 1986; 50: 1095-1098.
- [11] Bishop AR, Lomdahl PS. Nonlinear dynamics in driven, damped sine-Gordon systems.

- Physica D 1986; 18: 54-66.
- [12] Bishop AR, Forest MG, McLaughlin DW, Overman II EA. A quasi-periodic route to chaos in a near-integrable pde. Physica D 1986; 23: 293-328.
- [13] Rödelsperger F, Kivshar YS, Benner H. Reshaping-induced chaos suppression. Phys Rev E 1995; 51: 869-872.
- [14] Tereshko V, Shchekinova E. Resonant control of the Rössler system. Phys Rev E 1998; 58: 423-426.
- [15] Yakushevich LV. Nonlinear physics of DNA. Chichester: Wiley & Sons; 1998.
- [16] Kivshar YS, Malomed BA. Dynamics of solitons in near integrable systems. Rev Mod Phys 1989; 61: 763-915.

1. Figure Captions

Figure 1. Shape function $R(m)$ (see the text) versus the shape parameter m .

Figure 2. Theoretical (solid line, cf. Eq. (16)) and numerically obtained (dots) chaotic threshold amplitude, γ_{th} , versus shape parameter, m , for $\delta = 0.2$ and $T = 2\pi/0.6$.

Figure 3. Space-time evolution of $\Phi(x, t)$ for the sG system through four driving periods for $\delta = 0.2, \gamma = 0.79, T = 2\pi/0.6$, and shape parameters $m = 0, 0.5, 0.8, 0.82, 0.898 \simeq m_{th}$, and 0.99.

